

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

We Claim:

1. A pharmaceutical composition comprising:

- (a) a solubilized HIV protease inhibiting compound or a combination of solubilized HIV protease inhibiting compounds, or pharmaceutically acceptable salts thereof;
- (b) a pharmaceutically acceptable organic solvent which comprises a medium and/or long chain fatty acid or a mixture thereof, and ethanol or propylene glycol;
- (c) water; and
- (d) optionally, a pharmaceutically acceptable surfactant.

2. The composition according to Claim 1 wherein said HIV protease inhibiting compound is (2S,3S,5S)-5-(N-(N-((N-methyl-N-((2-isopropyl-4-thiazolyl)-(methyl)amino)carbonyl)-L-valinyl)amino-2-(N-((5-thiazolyl)methoxy-carbonyl)-amino)-1,6-diphenyl-3-hydroxyhexane (ritonavir).

3. The composition according to Claim 1 wherein said combination of HIV protease inhibiting compounds is (2S,3S,5S)-5-(N-(N-((N-methyl-N-((2-isopropyl-4-thiazolyl)-(methyl)amino)carbonyl)-L-valinyl)amino-2-(N-((5-thiazolyl)methoxy-carbonyl)-amino)-1,6-diphenyl-3-hydroxyhexane (ritonavir) and (2S,3S,5S)-2-(2,6-dimethylphenoxyacetyl)-amino-3-hydroxy-5-(2S-(1-tetrahydropyrimid-2-onyl)-3-methyl-

butanoyl) amino-1,6-diphenylhexane (ABT-378).

4. The composition according to Claim 1 wherein said HIV protease inhibiting compound or combination of HIV protease inhibiting compounds is selected from the group consisting of:

(2S,3S,5S)-5-(N-(N-(N-methyl-N-(2-isopropyl-4-thiazolyl)-methyl)amino)carbonyl)-L-valinyl)amino-2-(N-(5-thiazolyl)methoxy-carbonyl)-amino)-1,6-diphenyl-3-

10 hydroxyhexane (ritonavir);

2S,3S,5S)-5-(N-(N-(N-methyl-N-((2-isopropyl-4-thiazolyl)methyl)amino)carbonyl)amino-1,6-diphenyl-3-

hydroxyhexane (ritonavir) and (2S,3S,5S)-2-(2,6-dimethylphenoxyacetyl)-amino-3-hydroxy-5-(2S-(1-

15 tetrahydropyrimid-2-onyl)-3-methyl-

butanoyl) amino-1,6-diphenylhexane;

N- (2 (R) -hydroxy-1

(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-

20 pentaneamide (indinavir);

N-tert-butyl-decahydro-2-[2(R)-hydroxy-4-phenyl-3(S)-[[N-(2-quinolylcarbonyl)-L-asparaginyl]amino]butyl]-(4aS,8aS)-isoquinoline-3(S)-carboxamide (saquinavir);

5 (S) -Boc-amino-4 (S) -hydroxy-6-phenyl-2 (R) -

25 phenylmethylexanoyl- (L) -Val- (L) -Phe-morpholin-4-ylamide;

1 -Naphthoxyacetyl-beta-methylthio-Ala-(2S, 3S)-

3-amino-2-hydroxy-4-butanoyl 1,3-thiazolidine-4-t-butylamide;

5-isoquinolinoxyacetyl-beta-methylthio-Ala-(2S,3S)-3-amino-2-hydroxy-4-butanoyl-1,3-thiazolidine-4-t-butylamide;
[1S-[1R-(R-),2S*]]-N¹ [3-[[[(1,1 - dimethylethyl)amino]carbonyl] (2-methylpropyl)amino]-2-hydroxy-1 - (phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide;
VX-478;
DMP-323;
DMP-450;
10 AG1343 (nelfinavir);
BMS 186,318;
SC-55389a;
BILA 1096 BS; and
U-140690 (tipranavir),
15 or a pharmaceutically acceptable salt thereof.

5. The composition according to Claim 1 wherein said medium and/or long chain fatty acid is oleic acid.

20 6. The composition according to Claim 1 wherein said surfactant is Polyoxyl 35 castor oil (Cremophor EL[®]).

7. The composition according to Claim 1 wherein the solution is encapsulated into a hard gelatin capsule or a soft
25 gelatin capsule.

8. The composition of Claim 1 wherein the solvent comprises (1) a pharmaceutically acceptable long chain fatty acid in the amount of from about 40% to about 75% by weight of the total solution; (2) ethanol or
5 propylene glycol in the amount of from about 3% to about 12% by weight of the total solution; and (3) water in the amount of from about 0.4% to about 1.5% by weight of the total solution.

10 9. The composition of Claim 1 wherein the solvent comprises (1) oleic acid in the amount of from about 40% to about 75% by weight of the total solution; (2) ethanol or propylene glycol in the amount of from about 3% to about 12% by weight of the total solution; and
15 (3) water in the amount of from about 0.4% to about 1.5% by weight of the total solution.

2/19/03
20 10. The composition of Claim 9 wherein the HIV protease inhibiting compound is selected from the group consisting of:

2S,3S,5S)-5-(N-(N-((N-methyl-N-((2-isopropyl-4-thiazolyl)methyl)amino)carbonyl)amino-1,6-diphenyl-3-hydroxyhexane (ritonavir);

25 2S,3S,5S)-5-(N-(N-((N-methyl-N-((2-isopropyl-4-thiazolyl)methyl)amino)carbonyl)amino-1,6-diphenyl-3-hydroxyhexane (ritonavir) and (2S, 3S, 5S)-2-(2,6-Dimethylphenoxyacetyl)

amino-3-hydroxy-5-[2S-(1-tetrahydro-pyrimid-2-onyl)-3-methyl
butanoyl] amino-1,6-diphenylhexane;
N-(2(R)-hydroxy-1
(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-
5 pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pent
aneamide (indinavir);
N-tert-butyl-decahydro-2-[2(R)-hydroxy-4-phenyl-3(S)-[[N-(2-qu
inolylcarbonyl)-L-asparaginy] amino]butyl]-(4aS,8aS)-isoquinol
ine-3(S)-carboxamide (saquinavir);
10 5(S)-Boc-amino-4(S)-hydroxy-6-phenyl-2(R)-
phenylmethylhexanoyl-(L)-Val-(L)-Phe-morpholin-4-ylamide;
1-Naphthoxyacetyl-beta-methylthio-Ala-(2S, 3S)-
3-amino-2-hydroxy-4-butanoyl 1,3-thiazolidine-4-t-butylamide;
5-isoquinolinoxyacetyl-beta-methylthio-Ala-(2S,3S)-3-amino-2-h
15 ydroxy-4-butanoyl-1,3-thiazolidine-4-t-butylamide;
[1S-[1R-(R-),2S*]]-N¹ [3-[[[(1,1 -
dimethylethyl) amino] carbonyl] (2-methylpropyl) amino]-2-
hydroxy-1 -(phenylmethyl)propyl]-2-[(2-
quinolinylcarbonyl) amino]-butanediamide;
20 VX-478;
DMP-323;
DMP-450;
AG1343 (nelfinavir);
BMS 186,318;
25 SC-55389a;
BILA 1096 BS; and
U-140690 (tipranavir),

002250 1603/560

or a pharmaceutically acceptable salt thereof.

11. The composition of Claim 9 wherein the HIV protease inhibiting compound is ritonavir, (2S, 3S, 5S)-2-(2,6-dimethylphenoxyacetyl) amino-3-hydroxy-5-(2S-(1-tetrahydro-pyrimid-2-onyl)-3-methylbutanoyl) amino-1,6diphenylhexane, indinavir, saquinavir, nelfinavir, or VX-478.

12. The composition of Claim 1 wherein the HIV protease inhibiting compound is ritonavir or a combination of ritonavir and another HIV protease inhibiting compound.

13. The composition of Claim 12 wherein the solution is encapsulated in a soft elastic gelatin capsule (SEC).

14. The composition of Claim 1 which comprises:

(a) ritonavir in the amount of from about 1% to about 30% by weight of the total solution;

(b) a pharmaceutically acceptable organic solvent which comprises (i) oleic acid in the amount of from about 15% to about 99% by weight of the total solution and (2) ethanol in the amount of from about 3% to about 12% by weight of the total solution; and

(c) water in the amount of from about 0.4% to about 1.5% by weight of the total solution; and

5 (d) polyoxyl 35 castor oil in the amount of
from about 0% to about 20% by weight of the total
solution.

15. The composition of Claim/14 which comprises:

10 (a) ritonavir in the amount of from about 5%
to about 10% by weight of the total solution,

(b) a pharmaceutically acceptable organic solvent which comprises (1) oleic acid in the amount of from about 70% to about 75% by weight of the total solution; and (2) ethanol in the amount of from about 3% about 12% by weight of the total solution;

(c) water in the amount of from about 0.4% to about 1.5% by weight of the total solution; and

(d) polyoxyl 35 castor oil in the amount of
20 about 6% by weight of the total solution.

16. The composition of Claim 15 wherein the solution is encapsulated in a soft elastic gelatin capsule (SEC).

25

17. The composition of Claim 1 which comprises:

(a) ritonavir and ABT-378 in the amount of from about 1% to about 45% by weight of the total solution;

5

(b) a pharmaceutically acceptable organic solvent which comprises (i) oleic acid in the amount of from about 15% to about 99% by weight of the total solution and (2) propylene glycol in the amount of from about 1% to about 15% by weight of the total solution; and

10

(c) water in the amount of from about 0.4% to about 1.5% by weight of the total solution.

15

18. The composition of Claim 17 which comprises:

(a) ritonavir and ABT-378 in the amount of from about 1% to about 45% by weight of the total solution,

20

(b) a pharmaceutically acceptable organic solvent which comprises (1) oleic acid in the amount of from about 70% to about 75% by weight of the total solution; and (2) propylene glycol in the amount of from about 1% about 8% by weight of the total solution; and

25

(c) water in the amount of from about 0.4% to about 1.5% by weight of the total solution.

